



ALTABAX PRIOR AUTHORIZATION
SD DEPARTMENT OF SOCIAL SERVICES
MEDICAL SERVICES DIVISION

Fax Completed Form to:
866-254-0761
For questions regarding this
Prior authorization, call
866-705-5391

SD Medicaid requires that patients receiving a prescription for Altanax must first try and fail MUPIROCIN.

- Patients must use generic mupirocin for a minimum of 5 days for the trial to be considered a failure.
- Patients diagnosed with MRSA may be approved to use Altanax first-line.

Part I: RECIPIENT INFORMATION (To be completed by physician's representative or pharmacy):

RECIPIENT NAME:	RECIPIENT MEDICAID ID NUMBER:
Recipient Date of birth: / /	

Part II: PHYSICIAN INFORMATION (To be completed by physician's representative or pharmacy):

PHYSICIAN NAME:		PHYSICIAN PROVIDER NUMBER:	
City:	State:	PHONE: ()	FAX: ()

Part III: TO BE COMPLETED BY PHYSICIAN:

Requested Dosage: (must be completed)	Diagnosis for this request:
--	------------------------------------

Qualifications for coverage:

<input type="checkbox"/> Failed trial of mupirocin in the last 90 days	Was mupirocin trial for at least 5 days? <input type="checkbox"/> YES <input type="checkbox"/> NO
--	--

Adverse Reaction (attach FDA Medwatch form) or contraindication to mupirocin: (provide description below):

Medical Justification for use of Altanax without trial of mupirocin:

Physician Signature:	Date:
----------------------	-------

Part IV: PHARMACY INFORMATION

PHARMACY NAME:	SD MEDICAID PROVIDER NUMBER:
Phone: ():	FAX: ()
Drug:	NDC#:

Part V: FOR OFFICIAL USE ONLY

Date: / /	Initials: _____
Approved - Effective dates of PA: From: / /	To: / /
Denied: (Reasons)	